arXiv:1701.02974v1 [math.NT] 11 Jan 2017

A GENERALIZATION OF GAUSS' TRIANGULAR THEOREM

JANGWON JU AND BYEONG-KWEON OH

Abstract. A quadratic polynomial $\Phi_{a,b,c}(x,y,z) = x(ax+1) + y(by+1) + z(cz+1)$ is called universal if the diophantine equation $\Phi_{a,b,c}(x,y,z) = n$ has an integer solution x, y, z for any non negative integer n. In this article, we show that if $(a,b,c) = (2,2,6), (2,3,5)$ or $(2,3,7)$, then $\Phi_{a,b,c}(x,y,z)$ is universal. These were conjectured by Sun in [6].

1. introduction

A triangular number is a number represented as dots or pebbles arranged in the shape of an equilateral triangle. More precisely, the n-th triangular number is defined by $T_n = \frac{n(n+1)}{2}$ for any non negative integer n.

In 1796, Gauss proved that every positive integer can be expressed as a sum of three triangular numbers, which was first asserted by Fermat in 1638. This follows from the Gauss-Legendre theorem, which states that every positive integer which is not of the form $4^k(8l+7)$ with non negative integers k and l, is a sum of three squares of integers.

In general, a ternary sum $aT_x + bT_y + cT_z$ $(a,b,c > 0)$ of triangular numbers is called *universal* if for any non negative integer n, the diophantine equation

$$aT_x + bT_y + cT_z = n$$

has an integer solution x, y, z. In 1862, Liouville generalized the Gauss' triangular theorem by proving that a ternary sum $aT_x + bT_y + cT_z$ of triangular numbers is universal if and only if (a,b,c) is one of the following triples:

$$(1,1,1),\quad (1,1,2),\quad (1,1,4),\quad (1,1,5),\quad (1,2,2),\quad (1,2,3),\quad (1,2,4).$$

Recently, Sun in [6] gave an another generalization of the Gauss' triangular theorem. Since $\{T_x : x \in \mathbb{N} \cup \{0\}\} = \{T_x : x \in \mathbb{Z}\} = \{x(2x+1) : x \in \mathbb{Z}\}$, Gauss' triangular theorem implies that the equation $x(2x+1) + y(2y+1) + z(2z+1) = n$ has an integer solution x, y, z for any non negative integer n. Inspired by this motivation, he defined in [6] that for positive integers $a \leqslant b \leqslant c$, a ternary sum $\Phi_{a,b,c}(x,y,z) = x(ax+1) + y(by+1) + z(cz+1)$ is *universal* if the equation

$$x(ax+1) + y(by+1) + z(cz+1) = n$$

2000 *Mathematics Subject Classification.* Primary 11E12, 11E20.

Key words and phrases. Triangular theorem, Universal polynomials.

This work of the first author was supported by BK21 PLUS SNU Mathematical Sciences Division.

This work of the second author was supported by the National Research Foundation of Korea (NRF-2014R1A1A2056296).

has an integer solution for any non negative integer n. He showed that if $\Phi_{a,b,c}(x,y,z)$ is universal, then (a,b,c) is one of the following 17 triples:

$$\begin{array}{lllllll} (1,1,2), & (1,2,2), & (1,2,3), & (1,2,4), & (1,2,5), & (2,2,2), \\ (2,2,3), & (2,2,4), & (2,2,5), & (2,2,6), & (2,3,3), & (2,3,4), \\ (2,3,5), & (2,3,7), & (2,3,8), & (2,3,9), & (2,3,10) & \end{array}$$

and proved the universalities of some candidates. In fact, the universality of each ternary sum of triangular numbers was proved except the following 6 triples:

$$\textbf{(2,2,6)}, \quad \textbf{(2,3,5)}, \quad \textbf{(2,3,7)}, \quad (2,3,8), \quad (2,3,9), \quad (2,3,10). \tag{1.1}$$

He also conjectured that for each of these six triples, $\Phi_{a,b,c}(x,y,z)$ is universal.

In this article, we prove that if (a,b,c) is one of the triples written in boldface in the above candidates, then $\Phi_{a,b,c}(x,y,z)$ is, in fact, universal. To prove the universality, we use the method developed in [4]. We briefly review this method in Section 2 for those who are unfamiliar with it.

Let f be a positive definite integral ternary quadratic form. A symmetric matrix corresponding to f is denoted by M_f. For a diagonal form $f(x,y,z) = a_1x^2+a_2y^2+a_3z^2$, we simply write $M_f = \langle a_1, a_2, a_3\rangle$. The genus of f is the set of all integral forms that are locally isometric to f, which is denoted by $\mathrm{gen}(f)$. The set of all integers that are represented by the genus of f (f itself) is denoted by $Q(\mathrm{gen}(f))$ ($Q(f)$, respectively). For an integer a, we define

$$R(a,f) = \{(x,y,z) \in \mathbb{Z}^3 : f(x,y,z) = a\} \quad \text{and} \quad r(a,f) = |R(a,f)|.$$

For a set S, we define $\pm S = \{s : s \in S \text{ or } -s \in S\}$. For two $v = (v_1,v_2,v_3), v' = (v_1',v_2',v_3') \in \mathbb{Z}^3$ and a positive integer s, we write $v \equiv v' \pmod{s}$ if $v_i \equiv v_i' \pmod{s}$ for any $i \in \{1,2,3\}$.

Any unexplained notations and terminologies can be found in [2] or [5].

2. General tools

Let $a \leqslant b \leqslant c$ be positive integers. Recall that a ternary sum

$$\Phi_{a,b,c}(x,y,z) = x(ax+1) + y(by+1) + z(cz+1)$$

is said to be universal if the equation $\Phi_{a,b,c}(x,y,z) = n$ has an integer solution for any non negative integer n. Note that $\Phi_{a,b,c}(x,y,z)$ is universal if and only if the equation

$$bc(2ax+1)^2 + ac(2by+1)^2 + ab(2cz+1)^2 = 4abcn + bc + ac + ab =: \Psi_{a,b,c}(n)$$

has an integer solution for any non negative integer n. This is equivalent to the existence of an integer solution X, Y and Z of the diophantine equation

$$bcX^2 + acY^2 + abZ^2 = \Psi_{a,b,c}(n)$$

satisfying the following congruence condition

$$X \equiv 1 \pmod{2a},\ Y \equiv 1 \pmod{2b} \text{ and } Z \equiv 1 \pmod{2c}.$$

In some particular cases, representations of quadratic forms with some congruence condition corresponds to representations of a subform which is suitably taken (for details, see [1]).

For the representation of an arithmetic progression by a ternary quadratic form, we use the method developed in [1], [3] and [4]. We briefly introduce this method for those who are unfamiliar with it.

Let d be a positive integer and let a be a non negative integer ($a \leqslant d$). We define

$$S_{d,a} = \{dn + a \mid n \in \mathbb{N} \cup \{0\}\}.$$

For integral ternary quadratic forms f, g, we define

$$R(g, d, a) = \{v \in (\mathbb{Z}/d\mathbb{Z})^3 \mid vM_g v^t \equiv a \pmod{d}\}$$

and

$$R(f, g, d) = \{T \in M_3(\mathbb{Z}) \mid T^t M_f T = d^2 M_g\}.$$

A coset (or, a vector in the coset) $v \in R(g, d, a)$ is said to be *good* with respect to f, g, d and a if there is a $T \in R(f, g, d)$ such that $\frac{1}{d} \cdot vT^t \in \mathbb{Z}^3$. The set of all good vectors in $R(g, d, a)$ is denoted by $R_f(g, d, a)$. If every vector contained in $R(g, d, a)$ is good, we write $g \prec_{d,a} f$. If $g \prec_{d,a} f$, then by Lemma 2.2 of [3], we have

$$S_{d,a} \cap Q(g) \subset Q(f).$$

Note that the converse is not true in general.

Theorem 2.1. *Under the same notations given above, assume that there is a partition* $R(g, d, a) - R_f(g, d, a) = (P_1 \cup \cdots \cup P_k) \cup (\widetilde{P}_1 \cup \cdots \cup \widetilde{P}_{k'})$ *satisfying the following properties: for each* P_i*, there is a* $T_i \in M_3(\mathbb{Z})$ *such that*

(i) T_i *has an infinite order;*
(ii) $T_i^t M_g T_i = d^2 M_g$*;*
(iii) *for any vector* $v \in \mathbb{Z}^3$ *such that* $v \,(mod\ d) \in P_i$, $\frac{1}{d} \cdot vT_i^t \in \mathbb{Z}^3$ *and* $\frac{1}{d} \cdot vT_i^t \pmod{d} \in P_i \cup R_f(g, d, a)$,

and for each $\widetilde{P}_j$*, there is a* $\widetilde{T}_j \in M_3(\mathbb{Z})$ *such that*

(iv) $\widetilde{T}_j^t M_g \widetilde{T}_j = d^2 M_g$*;*
(v) *for any vector* $v \in \mathbb{Z}^3$ *such that* $v \,(mod\ d) \in \widetilde{P}_j$, $\frac{1}{d} \cdot v\widetilde{T}_j^t \in \mathbb{Z}^3$ *and* $\frac{1}{d} \cdot v\widetilde{T}_j^t \pmod{d} \in P_1 \cup \cdots \cup P_k \cup R_f(g, d, a)$.

Then we have

$$(S_{d,a} \cap Q(g)) - \bigcup_{i=1}^{k} g(z_i)\mathcal{S} \subset Q(f),$$

where the vector z_i *is a primitive eigenvector of* T_i *and* $\mathcal{S}$ *is the set of squares of integers.*

Proof. See Theorem 2.3 in [1]. □

3. Universality of $x(ax+1)+y(by+1)+z(cz+1)$

In this section, we prove that if $(a,b,c) = (2,2,6),(2,3,5)$ or $(2,3,7)$, then $\Phi_{a,b,c}(x,y,z)$ is universal. For the list of all candidates, see (1.1) in the introduction.

Lemma 3.1. *For any positive integer n, there are integers a,b and c such that $a^2+3b^2+3c^2 = 24n+7$ and $a \equiv b \equiv c \pmod 4$, whereas*

$$a \not\equiv b \pmod 8 \quad \text{or} \quad a \not\equiv c \pmod 8.$$

Proof. Since the class number of $f(x,y,z) = x^2+3(x-4y)^2+3(x-4z)^2$ is one and $S_{24,7} \subset Q(\mathrm{gen}(f))$, there is an integer solution $x^2+3y^2+3z^2 = 24n+7$ with $x \equiv y \equiv z \pmod 4$. If we define $g(x,y,z) = x^2+(x-8y)^2+3(x-8z)^2$, then it suffices to show that for any positive integer n,

$$r(24n+7,f) - r(24n+7,g) > 0.$$

Note that the genus of g consists of

$$M_g = \begin{pmatrix} 7 & -1 & 3 \\ -1 & 55 & 27 \\ 3 & 27 & 111 \end{pmatrix},\ M_2 = \begin{pmatrix} 15 & 6 & 3 \\ 6 & 28 & 14 \\ 3 & 14 & 103 \end{pmatrix} \text{ and } M_3 = \begin{pmatrix} 28 & 10 & 2 \\ 10 & 31 & -13 \\ 2 & -13 & 55 \end{pmatrix}.$$

Furthermore, one may easily show by using the Minkowski-Siegel formula that

$$r(24n+7,f) = r(24n+7,\mathrm{gen}(f)) = 4r(24n+7,\mathrm{gen}(g)),$$

where

$$r(24n+7,\mathrm{gen}(g)) = \frac{1}{4}r(24n+7,g) + \frac{1}{4}r(24n+7,M_2) + \frac{1}{2}r(24n+7,M_3).$$

Therefore we have

$$r(24n+7,f) - r(24n+7,g) = r(24n+7,M_2) + 2r(24n+7,M_3).$$

Hence it suffices to show that every integer of the form $24n+7$ that is represented by g is also represented by M_2 or M_3.

One may show by a direct computation that

$$R(g,8,7) - R_{M_3}(g,8,7) = \pm\{(1,0,0),(3,0,0)\}.$$

Let $P_1 = R(g,8,7) - R_{M_3}(g,8,7)$ and $T_1 = \begin{pmatrix} 8 & -4 & 0 \\ 0 & -2 & -12 \\ 0 & 6 & 4 \end{pmatrix}$. Since P_1 and T_1 satisfy all conditions in Theorem 2.1,

$$(S_{24,7} \cap Q(g)) - 7\mathcal{S} \subset Q(M_3).$$

Assume that $24n+7 = 7t^2$ for some positive integer t. Then t has a prime divisor p greater than 3. Since g represents 7, $24n+7 = 7t^2$ is represented by M_2 or M_3 by Lemma 2.4 of [1]. Therefore we have $r(24n+7,M_2) + 2r(24n+7,M_3) > 0$ for any positive integer n, which completes the proof. □

Theorem 3.2. *The ternary sum $x(2x+1)+y(2y+1)+z(6z+1)$ is universal.*

Proof. It suffices to show that for any positive integer n,

$$3(4x+1)^2+3(4y+1)^2+(12z+1)^2=24n+7$$

has an integer solution x,y,z. By Lemma 3.1, there is an integer a,b,c such that

$$a^2+3b^2+3c^2=24n+7,\quad a\equiv b\equiv c \pmod 4 \quad\text{and}\quad a\not\equiv b \pmod 8.$$

By changing signs suitably, we may assume that $a\equiv b\equiv c\equiv 1 \pmod 4$. If $a\equiv 1 \pmod 3$, then everything is trivial. Assume that $a\equiv 2 \pmod 3$. In this case, note that

$$\left(\frac{a-3b}{2}\right)^2+3\left(\frac{a+b}{2}\right)^2+3c^2=a^2+3b^2+3c^2=24n+7$$

and

$$\frac{a-3b}{2}\equiv 1 \pmod 3 \quad\text{and}\quad \frac{a-3b}{2}\equiv 1 \pmod 4.$$

The theorem follows directly from this. □

Theorem 3.3. *The ternary sum $x(2x+1)+y(3y+1)+z(5z+1)$ is universal.*

Proof. Let $f(x,y,z)=6x^2+10(2y+z)^2+15z^2$. We show that every integer of the form $40n+31$ for some positive integer n is represented by f. Note that the genus of f consists of

$$M_f=\langle 6\rangle\perp\begin{pmatrix}25&5\\5&25\end{pmatrix},\ M_2=\langle 1,30,120\rangle \text{ and } M_3=\langle 10,15,24\rangle.$$

Note also that $S_{40,31}\subset Q(\mathrm{gen}(f))$. By a direct computation, one may easily show that $M_3\prec_{40,31} M_2$. Hence it suffices to show that every integer of the form $40n+31$ that is represented by M_2 is also represented by f.

Note that $R(M_2,40,31)-R_f(M_2,40,31)$ consists of the followings:

$$\begin{array}{llll}\pm(1,\pm7,10), & \pm(1,\pm7,30), & \pm(1,\pm13,0), & \pm(1,\pm13,20),\\ \pm(9,\pm3,0), & \pm(9,\pm3,20), & \pm(9,\pm17,10), & \pm(9,\pm17,30),\\ \pm(11,\pm3,10), & \pm(11,\pm3,30), & \pm(11,\pm17,0), & \pm(11,\pm17,20),\\ \pm(19,\pm7,0), & \pm(19,\pm7,20), & \pm(19,\pm13,10), & \pm(19,\pm13,30).\end{array}$$

Now, we define a partition of $R(M_2,40,31)-R_f(M_2,40,31)$ as follows:

$$\begin{aligned}P_1&=\{(m_1,m_2,m_3)\mid 2m_1+m_2-m_3\equiv 0 \pmod 5\},\\ \widetilde{P}_1&=\{(m_1,m_2,m_3)\mid 3m_1+m_2-m_3\equiv 0 \pmod 5\},\end{aligned}$$

and we also define

$$T_1=\begin{pmatrix}-20&-180&120\\-6&22&12\\-1&-3&-38\end{pmatrix},\ \widetilde{T}_1=\begin{pmatrix}-20&180&-120\\-6&-22&-12\\-1&3&38\end{pmatrix}.$$

Then one may easily show that it satisfies all conditions in Theorem 2.1. Note that the eigenvector of T_1 is $(-3,1,0)$ and $120n+31$ is not of the form $39t^2$ for any positive integer t. Therefore for any non negative integer n, the equation $f(x,y,z)=120n+31$ has an integer solution $(x,y,z)=(a,b,c)$. Since $a\equiv 2b+c\equiv c\equiv 1 \pmod 2, 2b+c\not\equiv 0 \pmod 3$ and $a\equiv\pm 1 \pmod 5$, the equation

$$15(4x+1)^2+10(6y+1)^2+6(10z+1)^2=120n+31$$

has an integer solution for any non negative integer n. This completes the proof. $\square$

Theorem 3.4. *The ternary sum $x(2x+1)+y(3y+1)+z(7z+1)$ is universal.*

Proof. Let $f(x,y,z)=6x^2+14y^2+21z^2$. We show that every integer of the form $21n+20$ for some positive integer n is represented by f. Note that the genus of f consists of

$$M_f=\langle 6,14,21\rangle,\ M_2=\begin{pmatrix}5&1&2\\1&17&-8\\2&-8&26\end{pmatrix},\ \text{ and }\ M_3=\langle 3,14,42\rangle.$$

Note also that $S_{21,20}\subset Q(\mathrm{gen}(f))$. One may easily show by a direct computation that $M_3\prec_{21,20} M_2$. Hence it suffices to show that every integer of the form $21n+20$ that is represented by M_2 is also represented by f.

One may show by a direct computation that the set $R(M_2,21,20)-R_f(M_2,21,20)$ consists of 52 vectors, which is the union of the following sets:

$$\begin{aligned}
&P_1=\pm\{(0,0,2),(0,0,5)\}, && P_2=\pm\{(0,4,0),(0,10,0)\},\\
&P_3=\pm\{(3,2,0),(3,16,0)\}, && P_4=\pm\{(2,0,0),(5,0,0)\},\\
&\\
&\widetilde{P}_1=\pm\{(6,9,10),(6,9,17)\}, && \widetilde{P}_2=\pm\{(2,19,19),(5,16,16)\},\\
&\widetilde{P}_3=\pm\{(2,13,4),(5,1,10)\}, && \widetilde{P}_4=\pm\{(2,1,10),(5,13,4)\},\\
&\widetilde{P}_5=\pm\{(3,7,12),(3,14,12)\}, && \widetilde{P}_6=\pm\{(6,5,12),(6,19,12)\},\\
&\widetilde{P}_7=\pm\{(3,0,5),(3,0,19)\}, && \widetilde{P}_8=\pm\{(1,2,2),(8,16,16)\},\\
&\widetilde{P}_9=\pm\{(9,9,10),(9,9,17)\}.
\end{aligned}$$

We also define

$$\begin{aligned}
&T_1=\begin{pmatrix}-13&-24&-21\\-6&-3&21\\4&-12&21\end{pmatrix},\ T_2=\begin{pmatrix}-3&-40&16\\6&3&24\\-6&4&11\end{pmatrix},\quad T_3=\begin{pmatrix}-9&30&-39\\12&9&3\\6&-6&-9\end{pmatrix},\\
&T_4=\begin{pmatrix}-1&36&0\\-12&-9&0\\-2&-12&21\end{pmatrix},\quad T_5=\begin{pmatrix}-9&0&-45\\4&-21&13\\10&0&1\end{pmatrix},\ T_6=\begin{pmatrix}-1&-30&-15\\6&-9&27\\10&6&3\end{pmatrix},\\
&T_7=\begin{pmatrix}-9&24&12\\4&15&-24\\10&6&3\end{pmatrix},\quad T_8=\begin{pmatrix}-3&36&-3\\6&-9&27\\-6&-12&15\end{pmatrix},\quad T_9=\begin{pmatrix}-21&0&-21\\0&7&21\\0&-14&21\end{pmatrix}.
\end{aligned}$$

Note that for any $i=1,2,\dots,9$,

$$T_i^tM_2T_i=21^2M_2\quad\text{and}\quad(21^2\cdot T_i^{-1})^tM_2(21^2\cdot T_i^{-1})=21^2\cdot M_2.$$

Let $G=R_f(M_2,21,20)$, which is the set of good vectors. For any i, let Q_i be the set P_k or $\widetilde{P}_s$ for some k or s. Let U (and V) be an integral matrix T_k or $21^2T_s^{-1}$ for some k or s. For any vector $v\in\mathbb{Z}^3$ such that $v\pmod{21}\in Q_1$, if

$$\frac{1}{21}\cdot vU^t\in\mathbb{Z}^3\quad\text{and}\quad\frac{1}{21}\cdot vU^t\pmod{21}\in G\cup Q_{j_1}\cup\cdots\cup Q_{j_k},$$

then we write

$$Q_1\xrightarrow{U}G\cup Q_{j_1}\cup\cdots\cup Q_{j_k}.$$

We also use the notation

$$Q_1 \xrightarrow{U} G \cup Q_2 \cup B \xrightarrow{V} G \cup Q_{j_1} \cup \cdots \cup Q_{j_k},$$

if for any vector $v \in \mathbb{Z}^3$ such that $v \pmod{21} \in Q_1$, $\frac{1}{21} \cdot vU^t \in \mathbb{Z}^3$ and either $\frac{1}{21} \cdot vU^t \pmod{21} \in G \cup Q_2$ or

$$\frac{1}{21}\left(\frac{1}{21} \cdot vU^t\right)V^t \in \mathbb{Z}^3 \quad \text{and} \quad \frac{1}{21}\left(\frac{1}{21} \cdot vU^t\right)V^t \in G \cup Q_{j_1} \cup \cdots \cup Q_{j_k}.$$

Here B stands for the set of vectors of the form $\frac{1}{21} \cdot vU^t \pmod{21}$ that are not contained in $G \cup Q_2$.

Now, one may show by direct computations that

$$\begin{aligned}
&P_1 \xrightarrow{T_1} G \cup B \xrightarrow{T_2} G \cup \widetilde{P}_2, && \widetilde{P}_2 \xrightarrow{T_3} G \cup B \xrightarrow{T_4} G \cup P_1,\\
&P_2 \xrightarrow{T_5} G \cup P_2 \cup B \xrightarrow{T_6} G, && P_4 \xrightarrow{T_9} G \cup P_4,\\
&\widetilde{P}_1 \xrightarrow{T_1} G \cup B \xrightarrow{T_2} G, && \widetilde{P}_3 \xrightarrow{21^2T_1^{-1}} G \cup P_1 \cup \widetilde{P}_1,\\
&\widetilde{P}_4 \xrightarrow{21^2T_2^{-1}} G \cup \widetilde{P}_2 \cup \widetilde{P}_3, && \widetilde{P}_5 \xrightarrow{T_5} G \cup P_2 \cup B \xrightarrow{T_6} G,\\
&P_3 \xrightarrow{T_7} G \cup \widetilde{P}_3 \cup B \xrightarrow{T_8} G \cup P_3 \cup \widetilde{P}_5, && \widetilde{P}_6 \xrightarrow{21^2T_5^{-1}} G \cup P_2 \cup \widetilde{P}_5,\\
&\widetilde{P}_7 \xrightarrow{21^2T_8^{-1}} G \cup P_3 \cup \widetilde{P}_6, && \widetilde{P}_8 \xrightarrow{21^2T_9^{-1}} G \cup \widetilde{P}_7,\\
&\widetilde{P}_9 \xrightarrow{21^2T_8^{-1}} G \cup P_3 \cup \widetilde{P}_6.
\end{aligned}$$

As a sample of the above computations, assume that $v = (21x, 21y, 21z + 2)$ for some integers x, y, z. Note that $v \pmod{21} \in P_1$ and

$$w = \frac{1}{21} \cdot vT_1^t = (-13x - 24y - 21z - 2, -6x - 3y + 21z + 2, 4x - 12y + 21z + 2).$$

For any integers x, y, z, the vector $w \pmod{21}$ is contained in G or

$$w \pmod{21} = (16, 20, 11) \quad \text{or} \quad w \pmod{21} = (19, 2, 2).$$

If $w \pmod{21} = (16, 20, 11)$, then there are some integers a, b such that $(x, y, z) = (18 + 3a, a, b)$. Hence $w = (-236 - 63a - 21b, -106 - 21a + 21b, 74 + 21b)$. Now, one may show by a direct computation that

$$\frac{1}{21} \cdot wT_2^t \pmod{21} \in G \quad \text{or} \quad \frac{1}{21} \cdot wT_2^t \pmod{21} = (19, 2, 2) \in \widetilde{P}_2.$$

If $w \pmod{21} = (19, 2, 2)$, then there are some integers c, d such that $(x, y, z) = (3c, c, d)$. Hence $w = (-63c - 21d - 2, -21c + 21d + 2, 21d + 2)$. Now, one may show by a direct computation that

$$\frac{1}{21} \cdot wT_2^t \pmod{21} \in G \quad \text{or} \quad \frac{1}{21} \cdot wT_2^t \pmod{21} = (19, 2, 2) \in \widetilde{P}_2.$$

Note that the order of each $T_4T_3T_2T_1, T_5, T_8T_7$ and T_9 is infinite. Therefore, though the situation is slightly different from Theorem 2.1, we may still apply it to this case. The primitive integral eigenvectors of these four matrices are $(0, 0, 1), (0, 1, 0), (3, 2, 0)$ and $(1, 0, 0)$, respectively. Therefore if $168n + 41$ is not of the form $17t^2$ for some positive integer t, then $168n + 41$ is represented by f. Assume that $168n + 41 = 17t^2$ for some positive integer t. Then t has a prime divisor relatively prime to $2 \cdot 3 \cdot 7$. Since $M_2, M_3 \in \mathrm{spn}(f)$, and both M_2 and

M_3 represent 17, f represents $17t^2$ by Lemma 2.4 of [1]. Therefore the equation $f(x,y,z) = 168n + 41$ has an integer solution a, b, c for any non negative integer n. Since $a \equiv b \equiv c \equiv 1 \pmod 2$, $b \not\equiv 0 \pmod 3$ and $a \equiv \pm 1 \pmod 7$, the equation

$$21(4x+1)^2 + 14(6y+1)^2 + 6(14z+1)^2 = 168n + 41$$

has an integer solution for any non negative integer n. This completes the proof. □

References

[1] J. Ju, B.-K. Oh and B. Seo, *Ternary universal sums of generalized polygonal numbers*, submitted.

[2] Y. Kitaoka, *Arithmetic of quadratic forms*, Cambridge University Press, 1993.

[3] B.-K. Oh, *Regular positive ternary quadratic forms*, Acta Arith. **147**(2011), 233-243.

[4] B.-K. Oh, *Ternary universal sums of generalized pentagonal numbers*, J. Korean Math. Soc. **48**(2011), 837-847.

[5] O. T. O'Meara, *Introduction to quadratic forms*, Springer Verlag, New York, 1963.

[6] Z. W. Sun, *On* $x(ax+1) + y(by+1) + z(cz+1)$ *and* $x(ax+b) + y(ay+c) + z(az+d)$, J. Number Theory **171**(2017), 275-283.

Department of Mathematical Sciences, Seoul National University, Seoul 08826, Korea

E-mail address: jjw@snu.ac.kr

Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Seoul 08826, Korea

E-mail address: bkoh@snu.ac.kr